



Pearson

# Mark Scheme (Results)

Summer 2017

Pearson Edexcel International GCSE  
In Mathematics B (4MB0) Paper 01R

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2017

Publications Code 4MB0\_01R\_1706\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
  - M marks: method marks
  - A marks: accuracy marks
  - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
  - cao – correct answer only
  - ft – follow through
  - isw – ignore subsequent working
  - SC - special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - awrt – answer which rounds to
  - eeoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.


Q	Working	Answer	Mark	Notes
1 (a)		57900	1	B1
(b)		0.05	1	B1
				<b>Total 2 marks</b>

2	$3200000$ or $32 \times 10^5$		2	M1
		$3.2 \times 10^6$		A1
				<b>Total 2 marks</b>

3	$(3x =)7 - 2$		2	M1
		$5\frac{1}{3}$ (oe)		A1
				<b>NB do not allow 1.66</b>
				<b>Total 2 marks</b>

4	$14\frac{1}{30} \times 360$ (oe)		2	M1
		168		A1
				<b>Total 2 marks</b>

5	$\pm \frac{96-90}{96}$ or $\pm \frac{6}{96}$ or $\frac{90}{96} \times 100$		3	M1
	$\pm \frac{96-90}{96} \times 100$ or $100 - \frac{90}{96} \times 100$			M1 DEP
		$6\frac{1}{4}$ (oe)		A1
				$6.25, \frac{25}{4}$
	NB -6.25 score MIMIA0			<b>Total 3 marks</b>

6	(a) (i)		$x \geq -5$	2	B1
	(ii)		$x < 6$		B1
	(b)	Single Line -5		1	B1
					<b>Total 3 marks</b>

7		$\Delta = 5 \times -2 - 4 \times -3 (=2)$		3	M1
		$\frac{1}{\Delta} \begin{pmatrix} -2 & 3 \\ -4 & 5 \end{pmatrix}$			M1 (Indep) allow $\Delta=1$
		$\begin{pmatrix} -1 & 1\frac{1}{2} \\ -2 & 2\frac{1}{2} \end{pmatrix}$ (oe)			A1
					<b>Total 3 marks</b>



8		$2\sqrt{3} - 2\sqrt{2} + \sqrt{6} \times \sqrt{3} - \sqrt{6} \times \sqrt{2}$ or $2\sqrt{3} - 2\sqrt{2} + \sqrt{18} - \sqrt{12}$		3	M1
		$\sqrt{18} = 3\sqrt{2}$ and $\sqrt{12} = 2\sqrt{3}$ or $\sqrt{6} \times \sqrt{3} = 3\sqrt{2}$ and $\sqrt{6} \times \sqrt{2} = 2\sqrt{3}$			M1 (DEP)
		$2\sqrt{3} - 2\sqrt{2} + 3\sqrt{2} - 2\sqrt{3}$	$\sqrt{2}$		A1 dep on M2 CSO
					<b>Total 3 marks</b>

NB Answers derived from decimal expansions score M0M0A0

<b>9</b>	tangent drawn at (2, 2)			3	M1	Must only touch curve at (2,2)
	$\frac{y \text{ increment}}{x \text{ increment}}$				M1(DEP)	Attempt to find gradient for their tangent.
			3		A1	dep on tangent drawn ft their tangent providing $2.5 \leq \text{gradient} \leq 4$
<b>Total 3 marks</b>						

<b>10</b>	Shaded area = $(2r)^2 - \pi r^2$ , $0.858 r^2$ or better. OR $\frac{\pi(r^2)}{4(r^2)} \times 100$			3	M1	
	$\frac{4r^2 - \pi r^2}{4r^2} \times 100\%$ OR $100 - \frac{\pi(r^2)}{4(r^2)} \times 100$				M1 DEP	
			21.5		A1	awrt 21.4 or 21.5
<b>Total 3 marks</b>						

<b>11</b>	(a)			1	B1	
	(b)		$Q, Z, N$ correctly labelled $\{4, \sqrt{2}, -3, \frac{2}{3}\}$ four elements in correct position	2	B2	or B1 for two in correct position Do not allow repeated elements in different areas
<b>Total 3 marks</b>						

<b>12</b>			$\begin{pmatrix} 10 & 9 \\ 13 & 9 \\ -4 & -10 \end{pmatrix}$	3	M1 A1 A1	Correct dimension and at least one correct entry. Any three correct entries. All correct.
NB for a transcription error allow at most one value changed or two values transposed.						
<b>13</b>	(a)			1	B1	Correct region shaded.
	(b)			1	B1	Correct region shaded.
	(c)		$(F \cup G) \cap H'$ (oe)	1	B1	Accept $(F \cap H') \cup (G \cap H')$ or $(F \cap G' \cap H') \cup (F \cap G \cap H') \cup (F' \cap G \cap H')$ Do not accept $F \cup G \cap H'$
					<b>Total 3 marks</b>	



<b>14</b>	$\frac{6}{10} \times \frac{4}{10}$ or $\frac{4}{10} \times \frac{6}{9}$ or $\frac{6}{10} \times \frac{6}{10} + \frac{4}{10} \times \frac{3}{9}$		3	M1
	$\frac{6}{10} \times \frac{4}{10} + \frac{4}{10} \times \frac{6}{9}$ or $1 - \left( \frac{6}{10} \times \frac{6}{10} + \frac{4}{10} \times \frac{3}{9} \right)$			M1 DEP
		$\frac{38}{75}$ (oe)		A1 AWRT 0.507
				<b>Total 3 marks</b>

<b>15</b>	SF = $\sqrt[3]{\frac{1350}{400}}$ ( $= \frac{3}{2}$ ) or $\sqrt[3]{\frac{400}{1350}}$ ( $= \frac{2}{3}$ ) or $\sqrt[3]{1350}$ : $\sqrt[3]{400}$ or 3:2 (oe)		3	M1	Obtain linear scale factor. This mark not awarded unless cube root taken at some stage.
	$360 \times \left( \sqrt[3]{\frac{1350}{400}} \right)^2$ or $360 \times 2.25$ or $360 \div \left( \sqrt[3]{\frac{400}{1350}} \right)^2$ or $360 \div \frac{4}{9}$			M1(DEP)	
		810		A1	
				<b>Total 3 marks</b>	

<b>16</b>	$\frac{(2x-1)(x+5)}{3(2x-1)(2x+1)}$ or $\frac{(2x-1)(x+5)}{(2x-1)(6x+3)}$		3	M1	factorise numerator
				M1	factorise denominator
		$\frac{x+5}{3(2x+1)}$		A1	These marks are independent accept $\frac{x+5}{6x+3}$
				<b>Total 3 marks</b>	

<b>17</b>	(a)	$\frac{40}{32}$ or $75$ or $\frac{5}{4}$ or $1\frac{1}{4}$ or $1.25$		2	M1		
	(b)	$\frac{1.5 + 40 + 10}{22.5 + 75 + 37.5} \times 60 = \frac{51.5}{135}$ $\frac{1.5 + 40 + 10}{0.375 + "1.25" + 0.625} = \frac{51.5}{2.25} \text{ (oe)}$	1 hour 15 minutes	2	A1 M1 FT 75 or 1.25 from (a)		
					A1	awrt 22.9	<b>Total 4 marks</b>

<b>18</b>		$3x + y = 7$ $3x + 21y = -39$	$21x + 7y = 49$ $x + 7y = -13$	$y = 7 - 3x$ $x + 7(7 - 3x) = -13$	$x = -13 - 7y$ $3 \times (-13 - 7y) + y = 7$	4	M1 M1 DEP	Two equations with equal $x$ or $y$ coefficients, or correctly isolate $x$ or $y$ . Correct choice to add or subtract equations, or substitute $x$ or $y$ into other equation	Allow a single mistake across both M marks.
						$x = 3.1$ (oe) $y = -2.3$ (oe)		A1 A1	

<b>19</b>	$\sin 42 = \frac{5.2}{AC}$		4	M1	Any fully correct equation containing AC
	$AC = \frac{5.2}{\sin 42}$ (= 7.77 or better)			M1 DEP	Any fully correct expression for AC
	$\tan CAD = \frac{4.5}{\sin 7.77^\circ}$			M1 INDEP	ft AC > 0 Any fully correct expression for CAD or a trig ratio of CAD.
		30.1		A1	
					<b>Total 4 marks</b>

<b>20</b>	(a)	$2x - \frac{16}{x^2}$	2	B2	B1 for 2x, B1 for $-\frac{16}{x^2}$ or $-16x^{-2}$
	(b)	$2x - \frac{16}{x^2} = 0$ (oe)	2	M1	$\frac{dy}{dx} = 0$ , ft their $\frac{dy}{dx}$ as long as this is not the same as y
				A1	CSO
					<b>Total 4 marks</b>

<b>21</b>	$6 + 2x = 14$		4	M1	
		$x = 4$		A1	
	$3y - 3x = -27$ or $3y - 3 \times "4" = -27$			M1	
		$y = -5$		A1	
	NB award both M marks if $\begin{pmatrix} 6 + 2x \\ 3y - 3x \end{pmatrix} = \begin{pmatrix} 14 \\ -27 \end{pmatrix}$ seen.				<b>Total 4 marks</b>

<b>22</b>	$\text{Ext}_{t_2} = \frac{360}{12} (=30)$	$\text{Int}_{t_2} = \frac{(2 \times 12 - 4)90}{12}$ (= 150)	4	M1
	$\text{Ext}_t = 54 - "30" (= 24)$	$\text{Int}_t = 360 - 54 - "150"$ (= 156)		M1 INDEP
	$n = \frac{360}{"24"}$	$\frac{(2n - 4)90}{n} = "156" \text{ (oe)}$	15	M1 DEP A1
				<b>Total 4 marks</b>

<b>23</b>	(a)		1	B1
	(b)	$r^2 - 9 = 46(r + 3)$ $r - 3 - 46 = 0$ or $r - 3 = 46$ or $(r - 49)(r + 3) = 0$ or $(r =) \frac{-(-46) + \sqrt{(-46)^2 - 4 \times 1 \times -147}}{2 \times 1}$	3	M1 M1 (DEP)
			49	A1
		NB for factorisation in (b) allow any expression which gives 2 out of 3 correct in $r^2 - 46r - 147$ eg. $(r - 50)(r + 4)$		<b>Total 4 marks</b>

<b>24</b>	(a)	$0.2 + x + x + 0.1 + 0.18 + x + 0.1 = 1,$ $3x + 0.58 = 1 \text{ (oe)}$	2	M1
	(b)		0.14 (oe)	A1 $\frac{7}{50}$ ft for $x > 0$
	(c)	$60 \times 0.2$	2	M1
			12	A1
				<b>Total 5 marks</b>

<b>25</b>	(a)	$\frac{2 \times 2.5 + 5}{2.5 - 3} = \left(\frac{10}{-0.5}\right)$		2	M1	Fully correct substitution into function
			-20		A1	
	(b)	$xy - 3y = 2x + 5$ $x(y - 2) = 3y + 5$	$xy - 3x = 2y + 5$ $y(x - 2) = 3x + 5$	3	M1 M1 DEP A1	
			$\frac{3x+5}{x-2}$ (oe)			$3 + \frac{11}{x-2}$
<b>Total 5 marks</b>						

<b>26</b>	(a)	$360 - 255 - (180 - 135)$ or 60 correct on diagram		2	M1	SC Answer of 300 M1A0
			60		A1	
	(b)			1	B1	Accept 60(°), 060(°), N60(°)E
	(c)	"60" - (180 - 135) or 180 - (360 - 255) - "60" (oe) or 15 correct on diagram		2	M1	
			15		A1	Accept 15(°), 015(°), N15(°)E
<b>Total 5 marks</b>						

27	(a) (i)	equal intersecting arcs, centres $A$ and $C$		4	M1 A1 Any length.
	(ii)	arc centre $A$ to intersect $AC$ and $AB$ , and equal intersecting arcs from these points of intersection	perpendicular bisector drawn		M1
	(b)	Arc, centre $B$ , radius 5 cm	angle bisector drawn	2	A1 M1 Must extend from $A$ to $BC$ .
			region $R$ shaded		A1 FT FT their perpendicular bisector and angle bisector. The area must be bounded by at least both of these and the correct arc.
					<b>Total 6 marks</b>

28	(a) (i)	$2\pi r = \frac{216}{360} \times 2\pi \times 5x$ (oe)		4	M1
			3x		A1
	(ii)	$(5x)^2 = ("3x")^2 + h^2$ (oe)			M1
			4x		A1
	(b)	$\frac{1}{3}\pi("3x")^2 \times "4x" = \frac{375\pi}{2}$		3	M1
		$x^3 = \frac{375 \times 3}{2 \times 9 \times 4} \left( = \frac{125}{8} \right)$ (oe)			M1 DEP
			$2\frac{1}{2}$ (oe)		A1
					$\frac{5}{2}, 2.5$
					<b>Total 7 marks</b>

