



Mark Scheme (Results)

October 2023

Pearson Edexcel International Advanced Level
In Biology (WBI13) Paper 01
Unit 3: Practical Skills in Biology I

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2023

Question Paper Log Number P75617A

Publications Code WBI13_01_2310_MS

All the material in this publication is copyright

© Pearson Education Ltd 2023

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|----------------------------|-------------|
| 1ai | <p>An answer that includes the following points</p> <ul style="list-style-type: none"> • (using a light microscope find cell) under low power (1) • then view under high power (1) • calibrate eyepiece graticule / use of stage micrometer • count number of (eyepiece) graticule units over the cell (1) • convert eyepiece graticule units (to microns using calibration) (1) • measure diameter at different positions / orientations (1) | IGNORE electron microscope | expert 5 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 1a ii | <p>A drawing with the following features</p> <ul style="list-style-type: none"> • drawing using continuous lines showing correct shape of cell (1) • nucleus drawn with continuous lines in correct position and correct size (1) • any two correct labels (1) • suitable scale line shown (1) | <p>Lose this mark if draw cell wall</p> <p>membrane / cytoplasm / nucleus / mitochondria IGNORE cell wall, wrong labels as long as 2 are correct</p> <p>could be a scale line with suitable measurement / magnification (about 1000 if a 6 cm drawing, others pro-rata)</p> | 4 |

| Question Number | Answer | Additional Information | Marspapers.com |
|-----------------|---|---|----------------|
| 1aiii | <p>A calculation which includes the following steps</p> <ul style="list-style-type: none"> • conversion of onion cell length to micrometres / cheek cell to mm (1) • division of onion cell length by diameter of human cell and answer quoted as a ratio (1) | <p>= 200 (μm) OR 0.06 mm</p> <p>= $200 \div 60$ OR $0.2 \div 0.06$</p> <p>3.33 : 1 ACCEPT 3.3, 3.33</p> <p>ACCEPT 1:0.3</p> <p>NOT 1:3.33, this gets 1 mark</p> | 2 |


| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---------------------|------|
| 1bi | <p>The following correctly named:</p> <ul style="list-style-type: none"> X- Sclerenchyma, Y- Phloem / Sieve (tube element), Z- Xylem (cell / vessel) (2) <p>All three for 2 marks 1 or 2 for one mark.</p> | | 2 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|----------------------|------|
| 1bii | <ul style="list-style-type: none"> {phloem / sieve} cell / correct letter from 1bi | consider ecf from bi | 1 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|-------------------------------------|------|
| 2ai | <p>An answer that includes the following points</p> <ul style="list-style-type: none"> addition of (Benedict's / Fehling's) solution (1) heat / place in water bath (and observe colour change) (1) | Water bath must be heated or > 50 C | 2 |

| Question Number | Answer | Additional Guidance | Mark | | | | | | | | | | | | | | | | | | | | | |
|-------------------------|--|--|------------|--|------------------|------|-----|------|-------|------------|-------|------------------------|-------------|----------|-------------------------|--------------|-----|-------------------------|--------------|----------|----------------------|-------|------|---|
| 2a ii | | e.g. <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Appearance</th> <th>Glucose concentration / g dm^{-3}</th> <th>Glucose quantity</th> </tr> </thead> <tbody> <tr> <td>Blue</td> <td>0.0</td> <td>None</td> </tr> <tr> <td>Green</td> <td>0.1 to 0.4</td> <td>Trace</td> </tr> <tr> <td>Green with precipitate</td> <td>0.5 to 10.0</td> <td>Very low</td> </tr> <tr> <td>Yellow with precipitate</td> <td>10.1 to 15.0</td> <td>Low</td> </tr> <tr> <td>Orange with precipitate</td> <td>15.1 to 20.0</td> <td>Moderate</td> </tr> <tr> <td>Red with precipitate</td> <td>>20.0</td> <td>High</td> </tr> </tbody> </table> | Appearance | Glucose concentration / g dm^{-3} | Glucose quantity | Blue | 0.0 | None | Green | 0.1 to 0.4 | Trace | Green with precipitate | 0.5 to 10.0 | Very low | Yellow with precipitate | 10.1 to 15.0 | Low | Orange with precipitate | 15.1 to 20.0 | Moderate | Red with precipitate | >20.0 | High | 2 |
| Appearance | Glucose concentration / g dm^{-3} | Glucose quantity | | | | | | | | | | | | | | | | | | | | | | |
| Blue | 0.0 | None | | | | | | | | | | | | | | | | | | | | | | |
| Green | 0.1 to 0.4 | Trace | | | | | | | | | | | | | | | | | | | | | | |
| Green with precipitate | 0.5 to 10.0 | Very low | | | | | | | | | | | | | | | | | | | | | | |
| Yellow with precipitate | 10.1 to 15.0 | Low | | | | | | | | | | | | | | | | | | | | | | |
| Orange with precipitate | 15.1 to 20.0 | Moderate | | | | | | | | | | | | | | | | | | | | | | |
| Red with precipitate | >20.0 | High | | | | | | | | | | | | | | | | | | | | | | |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|---|------|
| 2b i | An answer that includes the following points: <ul style="list-style-type: none"> • describe how mean obtained (1) • describe how standard deviation obtained (1) | e.g substitute mean and number of observations into equation. | 2 |

| Question Number | Answer | Additional Guidance | www.dynamicpapers.com |
|-----------------|--|--|-----------------------|
| 2bii | <p>A graph with the following features</p> <ul style="list-style-type: none"> • L1 x label mean urine glucose concentration / au (1) • L2 y- mean blood glucose concentration / au) (1) • P1 plots correct on a linear scale on both axes (1) • P2 standard deviation plotted correctly (1) • suitable line of best fit drawn (1) |  <p>Mean Blood Glucose Concentration / au</p> <p>Mean Urine Glucose Concentration / au</p> | 5 |

IGNORE extrapolation

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| 2biii | <p>An answer that includes the following points:</p> <ul style="list-style-type: none"> • there is a correlation between the 2 variables / variables may depend on each other / as urine glucose goes up blood glucose goes up (1) • however, the {SDs / error bars} overlap (1) • <u>suggesting that differences not significant</u>(1) | <p>NOT vice versa to the last alternative</p> <p>NOT range bars</p> | 3 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| 2biv | <p>An calculation that includes the following steps :</p> <ul style="list-style-type: none"> • substitution of values into equation (1) • solving of equation (1) • correct rounding (1) | <p>e.g.</p> $y = 29.3 \times 12 + 154.1$ $= 505.7$ <p>rounded to 506</p> <p>allow ecf for $12 = 29.3 \times X + 154.1$ $= - 4.849$ (ignore sign) Rounded to 4.85</p> | 3 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|---------------------|------|
| 3ai | <p>An answer with the following :</p> <ul style="list-style-type: none"> • metaphase and anaphase | | 1 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 3aii | <p>An answer which includes the following points:</p> <p>Similarities</p> <ul style="list-style-type: none"> • cytokinesis (occurs) (1) • two (genetically) identical cells formed (1) <p>Differences</p> <ul style="list-style-type: none"> • cell plate formed in plants and not in animals / animal cell contracts but plant cell forms {vesicles / wall} in middle / plants cells remain connected by {plasmodesmata / pits}, none in animal cells. (1) | accept description, if implied under differences | 3 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|---|------|
| 3bi | <ul style="list-style-type: none"> lectin {treatment / added) / lectin concentration (1) | | 1 |
| 3bii | <p>An answer that includes the following points :</p> <ul style="list-style-type: none"> correct ref to safety issue (1) <p>and 4 of the following:</p> <ul style="list-style-type: none"> grow roots of one group in lectin, the other in water (1) obtain root tip (1) (root tips) placed in {(warm) acid / macerated / teased / described} (1 (1) (root tips) then placed in appropriate named stain (1) (root tip) squashed (1) cells dividing and {not dividing / interphase / total} counted under microscope (1) | <p>e.g. rinsing tips in water wearing gloves / goggles</p> <p>e.g. (acetic / ethanoic) orcein / toluidine blue / methylene blue</p> | 5 |

| Question Number | Answer | Additional Guidance | Mark | | | | | | | | | |
|---------------------|---|---|------|-------------------------------|---------------------------------------|---------|-----|----|---------------------|-----|----|---|
| 3biii | <p>A table with the following features :</p> <ul style="list-style-type: none"> • suitable table drawn (1) • all headings correct (1) • all data correctly entered (1) | <table border="1"> <thead> <tr> <th></th> <th>number of cells in Interphase</th> <th>number of cells in a stage of Mitosis</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>148</td> <td>24</td> </tr> <tr> <td>Treated with lectin</td> <td>160</td> <td>88</td> </tr> </tbody> </table> <p>table must have 2 rows and 2 columns minimum Do not allow .0 Extras lose mp 3</p> | | number of cells in Interphase | number of cells in a stage of Mitosis | Control | 148 | 24 | Treated with lectin | 160 | 88 | 3 |
| | number of cells in Interphase | number of cells in a stage of Mitosis | | | | | | | | | | |
| Control | 148 | 24 | | | | | | | | | | |
| Treated with lectin | 160 | 88 | | | | | | | | | | |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 3biv | <p>A calculation that includes the following steps :</p> <ul style="list-style-type: none"> • determine total number of cells in mitosis in lectin treatment and total number of cells observed (1) • calculate percentage (1) | <p>e.g.</p> $88 \text{ and } 160 + 88 = 248$ $\frac{(88 / 248) \times 100}{35.5} = 36$ | 2 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|------------------------|------|
| 3bv | <p>An answer that includes the following points :</p> <ul style="list-style-type: none"> percentage in mitosis is greater with lectins than without (1) but we have no idea of variability / difference may not be significant / no SDs (1) | IGNORE number of cells | 2 |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|----------------------|------|
| 3bvi | <p>An answer that includes the following points :</p> <ul style="list-style-type: none"> use of Chi (squared test) (1) because need to test {the significance of the difference between observed and expected results / association / frequency data} (1) | Allow ecf for t-test | 2 |

